

Editorial Comment

Do Neutrophils Mediate the Phenomenon of Stunned Myocardium?*

ROBERT A. KLONER, MD, PhD, FACC

Los Angeles, California

Brief periods of transient ischemia have been associated with prolonged postischemic dysfunction—the syndrome of stunned myocardium (1). In the canine model of 15 min of coronary artery occlusion followed by reperfusion, recovery of function may require hours to days despite the absence of tissue necrosis (2,3). Biochemical abnormalities have been observed in stunned myocardium: subendocardial adenosine triphosphate (ATP) levels are reduced for at least 3 days after reperfusion of a coronary artery subjected to 15 min of occlusion (4). Stunned myocardium has been observed in patients receiving thrombolytic therapy for acute evolving myocardial infarction, as well as in patients undergoing exercise stress testing (5).

Mechanisms of myocardial stunning. There have been several proposed mechanisms for the phenomenon of stunned myocardium, including altered calcium flux, oxygen free radical damage and abnormal energy utilization (6). Oxygen free radicals have been implicated as a cause of stunned myocardium by a number of investigators (7-9). These cytotoxic species such as superoxide anion and hydroxyl radical are known to be generated during ischemia and reperfusion (10). Oxygen free radicals recently have been documented in the coronary vein effluent of a canine model of 15 min of ischemia and reperfusion by utilizing electron paramagnetic resonance spectroscopy and a spin-trapping agent (10). In addition, oxygen radicals have been shown to inhibit the function of isolated sarcolemmal membrane preparations (11) and to depress function of isolated papillary muscle preparations (12). Several independent laboratories (7-9,10) including our own (8) have observed that administration of oxygen free radical scavengers, such as superoxide dismutase plus catalase, enhances the return of function of myocardium subjected to 15 min of ischemia

followed by reperfusion. This beneficial effect appears to occur without any improvement in ATP content of the tissue (8). Thus, oxygen free radicals do appear to play a role in the pathogenesis of stunned myocardium induced by brief periods of ischemia and reperfusion.

What is the origin of oxygen free radicals during ischemia and reperfusion? The xanthine oxidase reaction, neutrophils and electron transport chain of mitochondria are some of the potential sources (13,14). Perhaps the most controversial source associated with brief periods (15 min or less) of ischemia and reperfusion is the neutrophil. One problem with the concept that neutrophils are important in this setting is that their presence has not been described by histologic or electron microscopic examination in models of reversible ischemia (15). In fact, a recent indium-labeling study failed to document neutrophil influx into myocardium of dogs subjected to 12 min of ischemia and reperfusion (16).

Comparison with previous studies. Engler and Covell (17), however, showed that leukopak filtration of neutrophils from coronary reperfusate could blunt the phenomenon of stunned myocardium during brief periods of ischemia and reperfusion. In a preliminary report, Westlin and Mullane (18), also utilizing leukopak filtration, showed a transient improvement of stunned myocardium in neutropenic dogs. However, O'Neill et al. (19) showed that when neutrophils were suppressed with antiserum there was no effect on stunned myocardium. The present study by Jeremy and Becker (20) in this issue of the Journal used techniques similar to those of Engler and Covell (17) but the findings support the concept that neutrophils are not important in causing myocardial stunning in a model of brief coronary occlusion not associated with necrosis. The study is consistent with the data of O'Neill et al. (19) and with reported data that did not identify neutrophil infiltration into reversibly injured myocardium (15). The data are also consistent with models of isolated hearts that are known to develop stunning after global ischemia or hypoxia in which buffers devoid of neutrophils are used as the coronary perfusate but in which oxygen radical scavengers appear to be protective (21).

There are, however, some differences between the present study (20) and that of Engler and Covell (17); the latter suggested that very few neutrophils may be needed to cause stunning. Engler and Covell (17) therefore filtered neutrophils from blood entering both the left anterior descending and circumflex coronary arteries, whereas Jeremy and Becker (20) filtered only that blood entering the left anterior descending artery. Thus, in the study by Jeremy and Becker (20) neutrophils from the nonoccluded circumflex bed might have entered the bed of the left anterior descending coronary artery from collateral vessels, resulting in stunning. This occurrence would not explain the results of

*Editorials published in *Journal of the American College of Cardiology* reflect the views of the authors and do not necessarily represent the views of JACC or the American College of Cardiology.

From the Heart Institute, Hospital of the Good Samaritan and University of Southern California, Los Angeles, California.

Address for reprints: Robert A. Kloner, MD, Heart Institute of the Hospital of the Good Samaritan, 616 South Witmer Street, Los Angeles, California 90017.

O'Neill et al. (19) in which systemic neutropenia did not prevent stunning. It also would not explain a recent report by Schott et al. (22) in which myeloperoxidase activity (a marker of neutrophils) within stunned myocardium was similar to that of nonischemic controls. Furthermore, in that study, administration of antibody to the "adhesion-promoting" Mol glycoprotein on neutrophils did not prevent stunned myocardium.

The exact reasons for the differences in results among these studies is not clear. There may be as yet unknown or subtle factors that contribute to stunning. For example, the state of activation rather than the number of neutrophils present in the myocardium may be a crucial factor. Myers et al. (23) recently showed that neutrophils activated by phorbol myristate acetate depressed myocardial function and increased coronary vascular resistance in an isolated perfused rabbit heart model; unactivated neutrophils did not show this phenomenon. Also, R. Engler (personal communication, 1988) has suggested that other factors, such as the amount of vitamin E in diets of these animals, might help explain differences in results among laboratories.

Role of neutrophils. Although Jeremy and Becker's present study (20) shows that neutrophils do not play a role in causing stunning of the myocardium during brief periods of ischemia and reperfusion, neutrophils may play a more important role during prolonged periods (>1 h) of ischemia followed by reperfusion as might occur in the clinical situation of patients receiving thrombolytic therapy plus reperfusion for evolving acute myocardial infarction. Neutrophil infiltration has been observed on histologic (24,25), as well as by indium labeling early after reperfusion of irreversibly injured myocytes (26). We have observed zones of myocardium subjected to only a few hours of ischemia and reperfusion in which extensive neutrophil infiltration was present within the myocardium (24). Thus, while the early classic pathologic studies (27) suggested that neutrophil infiltration did not peak until 48 h after infarction, these studies were performed before the era of coronary reperfusion for the treatment of acute infarction. Coronary reperfusion may accelerate the early inflammatory response after infarction.

Engler et al. (28) have shown that the neutrophils may cause plugging of the microvasculature and contribute to the "no reflow" phenomenon in a model of prolonged occlusion (28). Some studies (25,29-31) have suggested that suppression of neutrophils in models of coronary occlusion with and without reperfusion may reduce myocardial infarct size and these studies have implicated the neutrophils as a cause of so-called reperfusion injury. Whether neutrophils contribute to stunned myocardium of peri-infarct areas in models of prolonged ischemia followed by reperfusion remains to be determined.

Finally, a word of caution. Neutrophils are a key element of the early inflammatory reaction that occurs after acute myocardial infarction. This early inflammatory reaction is

the first step in the healing process. Interventions that suppress neutrophils conceivably could alter the way in which myocardial infarcts heal. In fact, a host of anti-inflammatory agents, when administered shortly after coronary occlusion, have been shown to result in delayed healing, thin-walled scars and exacerbation of myocardial infarct expansion (32-34). However, one recent study (35) showed that administration of the oxygen free radical scavenger superoxide dismutase to reperfused infarcts did not result in excessive thinning of the infarcted ventricular wall.

Conclusions. The study by Jeremy and Becker (20) supports the concept that neutrophils do not play a role in postischemic ventricular dysfunction of myocardium that has been subjected to a brief period of ischemia and reperfusion not associated with necrosis. This study, however, does not rule out the possibility that neutrophils are important in models of more prolonged ischemia associated with irreversible cell injury. In fact, a preliminary study from the same research group (36) showed that perfusion of neutrophil-free blood after a 90 min period of ischemia reduced myocardial infarct size.

References

1. Braunwald E, Kloner RA. The stunned myocardium—prolonged, post-ischemic ventricular dysfunction. *Circulation* 1982;66:1146-9.
2. Heyndrickx GB, Baig H, Nellens P, Leusen I, Fishbein MC, Vatner SF. Depression of regional blood flow and wall thickening after brief coronary occlusions. *Am J Physiol* 1978;234:H653-H9.
3. Bolli R, Zhu WX, Thornby JI, O'Neill PG, Roberts R. Time course and determinants of recovery of function after reversible ischemia in conscious dogs. *Am J Physiol* 1988;254:H102-14.
4. DeBoer LWV, Ingwall JS, Kloner RA, Braunwald E. Prolonged derangement of canine myocardial purine metabolism following a brief coronary artery occlusion not associated with anatomic evidence of necrosis. *Proc Natl Acad Sci USA* 1980;77:5471-5.
5. Patel B, Kloner RA, Przyklenk K, Braunwald E. Postischemic myocardial "stunning": a clinically relevant phenomenon. *Ann Intern Med* 1988;108:627-9.
6. Kloner RA, Przyklenk K, Patel B. Altered myocardium states: the stunned and hibernating myocardium. *Am J Med* 1989;86(suppl 1A):14-22.
7. Myers ML, Bolli R, Lekich RF, Hartley CJ, Roberts R. Enhancement of recovery of myocardial function by oxygen free radical scavengers after reversible regional ischemia. *Circulation* 1985;72:915-21.
8. Przyklenk K, Kloner RA. Superoxide dismutase plus catalase improve contractile function in the canine model of "stunned myocardium". *Circ Res* 1986;58:148-56.
9. Gross GJ, Farber NE, Hardman HF, Warltier DC. Beneficial actions of superoxide dismutase and catalase in stunned myocardium of dogs. *Am J Physiol* 1986;250:H372-H7.
10. Bolli R, Patel BS, Jeroudi MO, Lai EK, McCay PB. Demonstration of free radical generation in "stunned" myocardium of intact dogs with the use of the spin trap L-Phenyl N-Tert-Butyl-Nitrone. *J Clin Invest* 1988;82:476-85.
11. Rowe GT, Manson NH, Caplan M, Hess ML. Hydrogen peroxide and hydroxyl radical mediation of activated leukocyte depression of cardiac sarcoplasmic reticulum, participation of the cyclooxygenase pathway. *Circ Res* 1983;53:584-91.

12. Blaustein AS, Schine L, Brooks WW, Fanburg BL, Bing OHL. Influence of exogenously generated oxidant species on myocardial function. *Am J Physiol* 1986;250:H595-9.
13. Bolli R. Oxygen-derived free radicals and postischemic myocardial dysfunction ("stunned myocardium"). *J Am Coll Cardiol* 1988;12:239-49.
14. Boveris A. Mitochondrial production of superoxide radical and hydrogen peroxide. *Adv Exp Med Biol* 1977;78:67-82.
15. Kloner RA, Ganote CE, Whalen D, Jennings RB. Effect of a transient period of ischemia on myocardial cells II. Fine structure during the first few minutes of reflow. *Am J Pathol* 1974;74:399-422.
16. Go LO, Murry CE, Richard VJ, Weischedel GR, Jennings RB, Reimer KA. Myocardial neutrophil accumulation during reperfusion after reversible or irreversible ischemic injury. *Am J Physiol* 1988;255:H1188-9.
17. Engler R, Covell JW. Granulocytes cause reperfusion ventricular dysfunction after 15 minutes of ischemia in the dog. *Circ Res* 1987;61:20-8.
18. Westlin W, Mullane KM. Leukocyte depletion attenuates postischemic myocardial dysfunction. *FASEB* 1988;2:A702.
19. O'Neill PG, Charlat ML, Michael LH, Roberts R, Bolli R. Influence of neutrophil depletion on myocardial function and flow after reversible ischemia. *Am J Physiol* 1989;256:H341-51.
20. Jeremy RW, Becker LC. Neutrophil depletion does not prevent myocardial dysfunction after brief coronary occlusion. *J Am Coll Cardiol* 1989;13:1155-63.
21. Shlafer M, Kane PF, Kirsh MM. Superoxide dismutase plus catalase enhance the efficacy of hypothermic cardioplegia to protect the globally ischemic, reperfused heart. *J Thorac Cardiovasc Surg* 1982;83:830-9.
22. Schott RJ, Nao BS, McClanahan TB, et al. Anti-Mol monoclonal antibody does not alter myocardial stunning. *Circulation* 1988;78(suppl II):II-78.
23. Myers ML, Webb CD, Moffat MP, Del Maestro RF. Activated neutrophils depress myocardial function in the perfused rabbit heart. *Circulation* 1988;78(suppl II):II-264.
24. Kloner RA, Alker KJ. Effect of streptokinase on intramyocardial hemorrhage, infarct size, and the no-reflow phenomenon during coronary reperfusion. *Circulation* 1984;70:513-21.
25. Mullane KM, Read N, Salmon JA, Moncada S. Role of leukocytes in acute myocardial infarction in anesthetized dogs: relationship to myocardial salvage by anti-inflammatory drugs. *J Pharmacol Exp Ther* 1984;228:510-22.
26. Engler RL, Dahlgren MD, Peterson MA, Dobbs A, Schmid-Schonbein GW. Accumulation of polymorphonuclear leukocytes during 3-h experimental myocardial ischemia. *Am J Physiol* 1986;251:H93-100.
27. Fishbein MC, Maclean D, Maroko PR. The histopathologic evolution of myocardial infarction. *Chest* 1978;73:843-4.
28. Engler RL, Schmid-Schonbein GW, Pavelec RS. Leukocyte capillary plugging in myocardial ischemia and reperfusion in the dog. *Am J Pathol* 1983;111:98-111.
29. Romson JL, Hook BG, Kunkel SL, et al. Reduction of the extent of ischemic myocardial injury by neutrophil depletion in the dog. *Circulation* 1983;67:1016-23.
30. Simpson PJ, Luccesi BR. Free radicals and myocardial ischemia and reperfusion injury. *J Lab Clin Med* 1987;110:13-30.
31. Simpson PJ, Mitsos SE, Ventura A, et al. Prostacyclin protects ischemic reperfused myocardium in the dog by inhibition of neutrophil activation. *Am Heart J* 1987;113:129-37.
32. Brown EJ Jr, Kloner RA, Schoen FJ, Hammerman H, Hale S, Braunwald E. Scar thinning due to ibuprofen administration following experimental myocardial infarction. *Am J Cardiol* 1983;51:877-83.
33. Hammerman H, Kloner RA, Schoen FJ, Brown EJ Jr, Hale S, Braunwald E. Indomethacin induced scar thinning following experimental myocardial infarction. *Circulation* 1983;67:1290-5.
34. Hammerman H, Kloner RA, Hale S, Schoen FJ, Braunwald E. Dose-dependent effects of short-term methylprednisolone on myocardial infarct extent, scar formation and ventricular function. *Circulation* 1983;68:446-52.
35. Werns SW, Shea MJ, Vaporciyan A, et al. Superoxide dismutase does not cause scar thinning after myocardial infarction. *J Am Coll Cardiol* 1987;9:898-902.
36. Litt M, Jeremy R, Weisman H, Becker L. Neutrophils cause reperfusion necrosis after ninety minutes myocardial ischemia (abstr). *Circulation* 1988;78(suppl II):II-16.